

## On $n$ -ary semigroups with adjoint neutral element

Wiesław A. Dudek and Vladimir V. Mukhin

### Abstract

We prove that we can adjoint an  $n$ -ary neutral element to an  $n$ -ary semigroup iff this semigroup is derived from a binary semigroup.

According to the general convention used in the theory of  $n$ -ary groupoids the sequence of elements  $x_i, x_{i+1}, \dots, x_j$  will be denoted by  $x_i^j$ . For  $j < i$  it is the empty symbol. If  $x_{i+1} = x_{i+2} = \dots = x_{i+t} = x$ , then instead of  $x_{i+1}^{i+t}$  we will write  $\overset{(t)}{x}$ . In this convention the symbol  $f(x_1, \dots, x_n)$  will be written as  $f(x_1^n)$ . Similarly, the symbol  $f(x_1^i, \overset{(t)}{x}, x_{i+t+1}^n)$  means  $f(x_1, \dots, x_i, \underbrace{x, \dots, x}_t, x_{i+t+1}, \dots, x_n)$ .

An  $n$ -ary groupoid  $(G, f)$  is called  $(i, j)$ -associative if

$$f(x_1^{i-1}, f(x_i^{n+i-1}), x_{n+i}^{2n-1}) = f(x_1^{j-1}, f(x_j^{n+j-1}), x_{n+j}^{2n-1}) \quad (1)$$

holds for all  $x_1, \dots, x_{2n-1} \in G$ . If this identity holds for all  $1 \leq i < j \leq n$ , then we say that the operation  $f$  is *associative* and  $(G, f)$  is called an  *$n$ -ary semigroup*. It is clear that an  $n$ -ary groupoid is associative if and only if it is  $(1, j)$ -associative for all  $j = 2, \dots, n$ . In the binary case (i.e. for  $n = 2$ ) it is a usual semigroup.

An  $n$ -ary semigroup  $(G, f)$  in which for all  $x_0, x_1, \dots, x_n \in G$  and all  $i \in \{1, \dots, n\}$  there exists an element  $z_i \in G$  such that

$$f(x_1^{i-1}, z, x_{i+1}^n) = x_0, \quad (2)$$

is called an  *$n$ -ary group*. It is clear that for  $n = 2$  we obtain a usual group.

Note by the way that in many papers  $n$ -ary semigroups ( $n$ -ary groups) are called  $n$ -semigroups ( $n$ -groups, respectively). Moreover, in many papers,

where the arity of the basic operation does not play a crucial role, is used the term *polyadic semigroups* (*polyadic groups*) (cf. [8]).

In the paper [1] written by W. Dörnte (under inspiration of Emmy Noether), he observed that any  $n$ -ary groupoid  $(G, f)$  of the form

$$f(x_1^n) = x_1 \circ x_2 \circ \dots \circ x_n \circ b,$$

where  $(G, \circ)$  is a group and  $b$  belongs to the center of this group, is an  $n$ -ary group but for every  $n > 2$  there are  $n$ -ary groups which are not of this form. In the first case we say that an  $n$ -ary groupoid  $(G, f)$  is *b-derived* (or *derived* if  $b$  is the identity of  $(G, \circ)$ ) from the group  $(G, \circ)$ , in the second – *irreducible*. Obviously, an  $n$ -ary operation derived from a binary associative operation is also associative in the above sense. An  $n$ -ary operation  $b$ -derived from an associative operation can be associative also in the case when  $b$  is not in the center. For example, the ternary operation  $b$ -derived from the multiplication of a nilpotent associative algebra of index 7 (the product of any 7 elements is 0) is trivially associative for every  $b$ .

In some  $n$ -ary groupoids there exists an element  $e$  (called an  *$n$ -ary neutral element*) such that

$$f(\overset{(i-1)}{e}, x, \overset{(n-i)}{e}) = x \quad (3)$$

holds for all  $x \in G$  and for all  $i = 1, \dots, n$ . There are  $n$ -ary semigroups (groups) with two, three and more neutral elements [9]. Also there are  $n$ -ary semigroups (groups too) in which all elements are neutral. All  $n$ -ary groups with this property are derived from the commutative group of the exponent  $k|(n-1)$  [2]. In  $n$ -ary group the set of neutral elements (if it is non-empty) forms an  $n$ -ary subgroup [5, 6]. In ternary groups each two neutral elements form a ternary subgroup. Other important properties of neutral elements one can find in [7] and [12].

As is it well known, to any semigroup  $(G, \circ)$  we can adjoin the identity  $e \notin G$  in this way that  $(G \cup \{e\}, \diamond)$  is a semigroup containing  $(G, \circ)$  as its semigroup. For this it is sufficient to define the operation  $\diamond$  as the extension of  $\circ$  putting  $x \diamond y = x \circ y$  for  $x, y \in G$ ,  $e \diamond e = e$  and  $x \diamond e = e \diamond x = x$  for  $x \in G$ .

Natural question is: *Is it possible to find the analogous construction for  $n$ -ary semigroups?* We prove below that the answer is positive.

First we characterize  $n$ -ary semigroups containing at least one  $n$ -ary neutral element.

**Lemma 1.** *An  $n$ -ary semigroup containing the neutral element is derived from a binary semigroup.*

*Proof.* Let  $e$  be the neutral element of an  $n$ -ary semigroup  $(G, f)$ . It is clear that  $(G, \circ)$ , where  $x \circ y = f(x, \overset{(n-2)}{e}, y)$ , is a semigroup and  $e$  is its neutral element. Direct computations shows that  $(G, f)$  is derived from  $(G, \circ)$ .  $\square$

From the above proposition we can deduce the following result firstly proved by W. Dörnte.

**Corollary 1.** *An  $n$ -ary group is derived from a binary group if and only if it has the neutral element.*

Note that any  $(i, j)$ -associative  $n$ -ary groupoid  $(G, f)$  with the neutral element in the center is an  $n$ -ary semigroup [3, 10, 11]. Such groupoid is associative also in the case when in the center of  $(G, f)$  lies at least one *neutral polyad (sequence)*, i.e., the sequence of elements  $a_2^n \in G$  such that  $f(x, a_2^n) = f(a_2^n, x) = x$  holds for all  $x \in G$  [3, 11]. Neutral sequences are in all  $n$ -ary groups ([8]), but not in all  $n$ -ary semigroups.

**Lemma 2.** *An  $n$ -ary semigroup derived from a binary semigroup possess a neutral sequence if and only if it contains the neutral element.*

*Proof.* Let  $(G, f)$  be derived from a semigroup  $(G, \circ)$ . If  $a_2^n$  is a neutral sequence of  $(G, f)$ , then  $e = a_2 \circ a_3 \circ \dots \circ a_n$  belongs to  $G$  and  $x \circ e = x \circ a_2 \circ a_3 \circ \dots \circ a_n = f(x, a_2^n) = x$  for all  $x \in G$ . Similarly  $e \circ x = x$ . This means that  $e$  is the identity of  $(G, \circ)$ . Hence it is the neutral element of an  $n$ -ary semigroup derived from  $(G, \circ)$ .

The converse statement is obvious.  $\square$

**Corollary 2.** *If an  $n$ -ary semigroup without neutral elements is derived from a binary semigroup then it does not possess any neutral sequence.*

**Proposition 1.** *A neutral element can be adjoint to any  $n$ -ary semigroup derived from a binary semigroup.*

*Proof.* Let  $n$ -ary semigroup  $(G, f)$  be derived from a binary semigroup  $(G, \circ)$ . Then to  $(G, \circ)$  we can add the identity  $e \notin G$  in this way that  $(G \cup \{e\}, \circ)$  becomes a semigroup with  $(G, \circ)$  as its subsemigroup. In an  $n$ -ary semigroup  $(G \cup \{e\}, g)$  derived from  $(G \cup \{e\}, \circ)$  the element  $e$  is neutral and  $f(x_1^n) = g(x_1^n)$  for  $x_1^n \in G$ . So, to  $(G, f)$  we can adjoint the neutral element  $e \notin G$ .  $\square$

**Proposition 2.** *If an  $n$ -ary semigroup  $(G, f)$  do not contains any neutral elements, then to  $(G, f)$  we can adjoint the neutral element if and only if  $(G, f)$  is derived from a binary semigroup.*

*Proof.* If to an  $n$ -ary semigroup  $(G, f)$  we can adjoin the neutral element  $e \notin G$ , then on  $G \cup \{e\}$  we can define the  $n$ -ary operation  $g$  such that  $g(x_1^n) = f(x_1^n)$  for all  $x_1^n \in G$ . By Lemma 1, an  $n$ -ary semigroup  $(G \cup \{e\}, g)$  is derived from the semigroup  $(G \cup \{e\}, *)$ , where  $x * y = g(x, \overset{(n-2)}{e}, y)$ . Obviously  $(G, *)$  is a subsemigroup of  $(G \cup \{e\}, *)$ . If not, then there are  $a, b \in G$  such that  $e = a * b$  which contradicts to the assumption on  $e$ . This means that  $(G, f)$  is an  $n$ -ary subsemigroup of  $(G \cup \{e\}, g)$  and it is derived from  $(G, *)$ .

The converse statement follows from Proposition 1.  $\square$

As a consequence of the above two propositions we obtain the following

**Theorem 1.** *To an  $n$ -ary semigroup  $(G, f)$  we can adjoin the neutral element if and only if  $(G, f)$  is derived from a binary semigroup.*

From the above proofs it follows that in an  $n$ -ary semigroup  $(G, f)$  derived from a semigroup  $(G, \circ)$  the adjoint  $n$ -ary neutral element is the adjoint identity of  $(G, \circ)$ .

**Theorem 2.** *For every  $n > 2$  there exists at least one  $n$ -ary semigroup (group) to which any  $n$ -ary neutral element cannot be adjoint.*

*Proof.* It is sufficient to prove that for every  $n > 2$  there exists at least one  $n$ -ary group without neutral elements.

At first consider the multiplicative group  $G = T(3, \mathbb{K})$  of triangular matrices of the form  $\begin{bmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{bmatrix}$ , where  $\mathbb{K}$  is a field of non-zero characteristic  $p$ . Then the map

$$\theta \begin{bmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & \alpha x & y \\ 0 & 1 & \beta z \\ 0 & 0 & 1 \end{bmatrix},$$

where  $\alpha$  is a primitive root of unity of degree  $n - 1$  and  $\alpha\beta = 1$ , is an automorphism of this group. It is not difficult to verify that the set  $G$  with the operation

$$f(A_1, A_2, \dots, A_n) = A_1 \cdot \theta(A_2) \cdot \theta^2(A_3) \cdot \dots \cdot \theta^{n-1}(A_n) \cdot B, \quad (4)$$

where  $B = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ , is an  $n$ -ary group.

This group do not contains any  $n$ -ary neutral element. Indeed, if  $A$  is its  $n$ -ary neutral element, then we have  $f(X, A, A, \dots, A) = f(A, X, A, \dots, A)$  for all  $X \in G$ . Whence, according to (4), we conclude  $X \cdot \theta A = A \cdot \theta X$ . Taking the identity matrix as  $X$ , we get  $\theta A = A$ . This proves that the matrix  $A$  belongs to the center of the group  $(G, \cdot)$ . Thus  $X \cdot A = A \cdot \theta X = \theta X \cdot A$ , which implies  $\theta X = X$  for all  $X \in G$ . This is not true. So,  $(G, f)$  is an  $n$ -ary group without neutral elements.

Now we give the another example of  $n$ -ary group without  $n$ -ary neutral elements.

Let  $\mathbb{C}$  be the set of complex numbers and let  $\omega$  be the primitive  $(n-1)$ -th root of unity. Then  $G = \mathbb{C}^3$  with the operation

$$\mathbf{x} \bullet \mathbf{y} = (x_1, x_2, x_3) \bullet (y_1, y_2, y_3) = (x_1 + y_1, x_2 + y_2 + x_1 y_3, x_3 + y_3)$$

is a group and  $\theta(x_1, x_2, x_3) = (\omega x_1, \omega^2 x_2, \omega x_3)$  is its automorphism.

It is not difficult to verify that  $(G, g)$ , where

$$g(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \mathbf{x}_1 \bullet \theta(\mathbf{x}_2) \bullet \theta^2(\mathbf{x}_3) \bullet \dots \bullet \theta^{n-1}(\mathbf{x}_n),$$

is an  $n$ -ary group. It is isomorphic to an  $n$ -ary group of triangular matrices from the proof of Theorem 3 in [4].

Similarly as in the previous case we can prove that  $(G, g)$  is not derived from any binary group.  $\square$

## References

- [1] **W. Dörnte**: *Untersuchungen über einen verallgemeinerten Gruppenbegriff*, Math. Z. **29** (1928), 1 – 19.
- [2] **W. A. Dudek**: *Autodistributive  $n$ -groups*, Commentationes Math. Annales Soc. Math. Polonae, Prace Matematyczne **23** (1983), 1 – 11.
- [3] **W. A. Dudek**: *On  $(i, j)$ -associative  $n$ -groupoids with the non-empty center*, Ricerche di Matematica (Napoli) **35** (1986), 105 – 111.
- [4] **W. A. Dudek**: *On the class of weakly semiabelian polyadic groups*, Discrete Math. Appl. **6** (1996), 427 – 433 (translation from Diskretn. Mat. **8** (1996), 40 – 46).
- [5] **W. A. Dudek**: *Idempotents in  $n$ -ary semigroups*, Southeast Asian Bull. Math. **25** (2001), 97 – 104.
- [6] **A. M. Gal'mak**:  *$n$ -ary subgroups of identities*, (Russian), Vesti Nats. Akad. Navuk Belarussi, ser. Fiz.-mat. **2** (2003), 25 – 30.

- [7] **M. S. Pop:** *On some relations on  $n$ -monoids*, Carpatian J. Math. **20** (2004), 87 – 94.
- [8] **E. L. Post:** *Polyadic groups*, Trans. Amer. Math. Soc. **48** (1940), 208 – 350.
- [9] **D. W. Robinson:**  *$n$ -groups with identity elements*, Math. Mag. **31** (1958), 255 – 258.
- [10] **B. Trpenovski:** *On  $n$ -groupoids containing central neutral elements*, (Makedonian), Bull. Soc. Math. et Phys. R. P. Macédoine **14** (1963), 31 – 39.
- [11] **B. Trpenovski and G. Čupona:** *Finitary associative operations with neutral elements*, (Makedonian), Bull. Soc. Math. et Phys. R. P. Macédoine **12** (1961), 15 – 24.
- [12] **D. Zupnik:** *Polyadic semigroups*, Publ. Math. (Debrecen), **14** (1967), 273 – 279.

Received December 28, 2005

W. A. Dudek

Institute of Mathematics and Computer Science, Wrocław University of Technology,  
50-370 Wrocław, Poland

E-mail: dudek@im.pwr.wroc.pl

V. V. Mukhin

Department of Mathematics, Cherepovets State University, 162 600 Cherepovets,  
Russia

E-mail: mukhin@chsu.ru